

Before using this product read the label carefully



# Maxforce<sup>®</sup> Ant

## Bait Granules

Reg. No. L5658 Act No. 36 of 1947  
Namibian Reg. No. N-AR 0591

### EXPERT



**KILLS THE COLONY  
NOT JUST  
FORAGING ANTS**

Ready-to-use granular bait for the control of **ANTS** (including **BROWN HOUSE ANT** and **BLACK SUGAR ANT**) outdoors around domestic premises and buildings, on lawns, turfgrass and other areas such as parks and golf courses.

**ACTIVE INGREDIENT:** Hydramethylinon.....10 g/kg

**REGISTRATION HOLDER:**

Bayer (Pty.) Ltd. Reg. No. 1968/011192/07  
P.O. Box 143 ISANDO 1600 SOUTH AFRICA TEL: (011) 921 5911  
MAXFORCE<sup>®</sup> is a registered trademark of the Bayer Group.

**225 g**



**CAUTION**



**WARNINGS:** Harmful if swallowed. May irritate skin and eyes. Toxic to fish. Keep out of reach of children, uninformed persons and animals. Store in a cool, dry place, away from food, feedstuffs and other insecticides. Keep container closed when not in use.

**PRECAUTIONS:** Avoid contact with skin or eyes. Wear gloves and wash hands thoroughly after use. Avoid contamination with other insecticides. Destroy empty container and do not re-use for any other purpose.

**DIRECTIONS FOR USE:** USE ONLY AS DIRECTED.

**For outdoor use only.**

The delayed action of the active ingredient allows foraging ants to bring the bait back to the nest so that the entire colony, including the queen, is destroyed. When properly used, this bait will control many common structure-invading ants before they enter a building. Do not apply other insecticides e.g. aerosols, sprays, etc. on top of or in close proximity to **Maxforce<sup>®</sup> Ant Bait Granules**, as it will reduce the effectiveness. Reduced activity and dead ants around the entrance of the nest should be noted around 7 days after baiting.

**1. Perimeter treatment**

Lightly sprinkle granules at the rate of 6 g/m<sup>2</sup> in a band 0,5 m wide around the perimeter of buildings where ants are active. Treat ant trails, cracks and crevices where ants may enter structures.

**2. Broadcast application**

Broadcast granules uniformly on lawns and other areas at the rate of 20 g/100 m<sup>2</sup>. Bait will remain effective, even after exposure to rain and sprinklers, for up to 7 days in direct sunlight.

**3. Ant nests**

Sprinkle approximately 30 g bait around active ant nests. Do not apply bait directly in or on the mound. If the mound is disturbed, ants will become agitated and may not feed on the bait.

For any product enquiry, contact Bayer - 011 921 5911. For poisoning emergencies only, Griffon Poison Information Centre may also be contacted - 082 446 8946.

© Bayer (Pty.) Ltd. All unauthorized reproduction of this label is forbidden.

[www.bayerhome.co.za](http://www.bayerhome.co.za)


Batch no.: See on container

Date manufactured: See on container

**For Scatter  
Treatment  
& Ant Nest  
Treatment:**



**For Perimeter  
Treatment:**



7998231D / 0712